



Year: 2017

Cold thermal irrigation decreases the ipsilateral gain of the vestibulo-ocular reflex

Tamás, L T ; Weber, K P ; Bockisch, C J ; Straumann, D ; Lasker, D M ; Büki, B ; Tarnutzer, A A

Abstract: OBJECTIVES: During head rotations, neuronal firing rates increase in ipsilateral and decrease in contralateral vestibular afferents. At low accelerations, this "push-pull mechanism" is linear. At high accelerations, however, the change of firing rates is nonlinear in that the ipsilateral increase of firing rate is larger than the contralateral decrease. This mechanism of stronger ipsilateral excitation than contralateral inhibition during high-acceleration head rotation, known as Ewald's second law, is implemented within the nonlinear pathways. The authors asked whether caloric stimulation could provide an acceleration signal high enough to influence the contribution of the nonlinear pathway to the rotational vestibulo-ocular reflex gain (rVOR gain) during head impulses. DESIGN: Caloric warm (44°C) and cold (24, 27, and 30°C) water irrigations of the left ear were performed in 7 healthy human subjects with the lateral semicircular canals oriented approximately earth-vertical (head inclined 30° from supine) and earth-horizontal (head inclined 30° from upright). RESULTS: With the lateral semicircular canal oriented earth-vertical, the strongest cold caloric stimulus (24°C) significantly decreased the rVOR gain during ipsilateral head impulses, while all other irrigations, irrespective of head position, had no significant effect on rVOR gains during head impulses to either side. CONCLUSIONS: Strong caloric irrigation, which can only be achieved with cold water, reduces the rVOR gain during ipsilateral head impulses and thus demonstrates Ewald's second law in healthy subjects. This unilateral gain reduction suggests that cold-water caloric irritation shifts the set point of the nonlinear relation between head acceleration and the vestibular firing rate toward a less acceleration-sensitive zone.

DOI: <https://doi.org/10.1097/AUD.0000000000000398>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-135798>

Journal Article

Accepted Version

Originally published at:

Tamás, L T; Weber, K P; Bockisch, C J; Straumann, D; Lasker, D M; Büki, B; Tarnutzer, A A (2017). Cold thermal irrigation decreases the ipsilateral gain of the vestibulo-ocular reflex. *Ear and Hearing*, 38(3):e193-e199.

DOI: <https://doi.org/10.1097/AUD.0000000000000398>

Cold thermal irrigation decreases the ipsilateral gain of the vestibulo-ocular reflex

L. T. Tamás MD¹, K. P. Weber MD^{2,3}, Ch. J. Bockisch PhD^{2,3,4}, D. Straumann MD², D. M. Lasker PhD⁵, B. Büki MD⁶, A. A. Tarnutzer MD²

¹Department of Otolaryngology, Petz Aladár County Hospital, Győr, Hungary

²Department of Neurology, University Hospital Zurich, University of Zurich, CH-8091 Zurich, Switzerland

³Department of Ophthalmology, University Hospital Zurich, University of Zurich, CH-8091 Zurich, Switzerland

⁴Department of Otorhinolaryngology, University Hospital Zurich, University of Zurich, CH-8091 Zurich, Switzerland

⁵Department of Otolaryngology - Head and Neck Surgery, Johns Hopkins University School of Medicine, Baltimore, 21287 MD, USA

⁶Department of Otolaryngology, Karl Landsteiner University Hospital Krems, 3500 Krems, Austria

Running title: Caloric irrigation and the head-impulse test

Corresponding author:

Alexander A. Tarnutzer, MD

Department of Neurology, University Hospital Zurich, Frauenklinikstr. 26
8091 Zurich, Switzerland

Phone: +41 44 255 11 11; Fax: +41 44 255 43 80

Email: alexander.tarnutzer@access.uzh.ch

Statistics:

Character count of the title (including spaces): 119

Word count of the abstract (max. 250): 241

Word count of the text (excluding figure legends, tables and references): 3245

Number of figures: 5

Conflict of interest and sources of funding:

Dr. Tamás, Dr. Bockisch, Dr. Lasker, Dr. Büki, Prof. Straumann and Dr. Tarnutzer report no conflict of interest. Dr. Weber acts as an unpaid consultant and has received funding for travel from GN Otometrics.

This study was supported by Dr. Dabbous Foundation, University of Zurich, Switzerland; Swiss National Science Foundation; and Betty and David Koetser Foundation, Zurich, Switzerland.

ABSTRACT

Objectives: During head rotations, neuronal firing rates increase in ipsilateral and decrease in contralateral vestibular afferents. At low accelerations, this ‘push-pull mechanism’ is linear. At high accelerations, however, the change of firing rates is non-linear in that the ipsilateral increase of firing rate is larger than the contralateral decrease. This mechanism of stronger ipsilateral excitation than contralateral inhibition during high-acceleration head rotation, known as Ewald’s second law, is implemented within the non-linear pathways. We asked whether caloric stimulation could provide an acceleration signal high enough to influence the contribution of the non-linear pathway to the rotational VOR-gain (rVOR-gain) during head impulses.

Design: Caloric warm (44°C) and cold (24, 27, 30°C) water irrigations of the left ear were performed in seven healthy human subjects with the lateral semicircular canals oriented approximately earth-vertical (head inclined 30° from supine) and earth-horizontal (head inclined 30° from upright).

Results: With the lateral semicircular canal oriented earth-vertical, the strongest cold caloric stimulus (24°C) significantly decreased the rVOR-gain during ipsilateral head impulses, while all other irrigations, irrespective of head position, had no significant effect on rVOR-gains during head impulses to either side.

Conclusions: Strong caloric irrigation, which can only be achieved with cold water, reduces the rVOR-gain during ipsilateral head impulses and thus demonstrates Ewald’s second law in healthy subjects. This unilateral gain-reduction suggests that cold-water caloric irritation shifts the set point of the non-linear relation between head acceleration and the vestibular firing-rate toward a less acceleration-sensitive zone.

INTRODUCTION

The six semicircular canals (SCCs) of the two labyrinths measure angular accelerations of the head about any axis in three-dimensional space. Within each cupula, this information is transformed into neuronal signals that are sent to the brainstem along the vestibular nerve. A major projection of vestibular signals feeds to a network that drives the rotational vestibulo-ocular reflex (rVOR). The rVOR ensures clear vision by eliciting compensatory eye movements in the opposite direction of head rotation with approximately the same angular velocity as the head. In normal-life conditions, head accelerations often occur with high frequencies, i.e. greater than 1 Hz, e.g. during locomotion (Grossman et al. 1988). To ensure a robust performance of the rVOR, elements of the SCCs and their afferents are specialized in transmitting information about head rotations of high angular acceleration to the vestibular nuclei (Fernandez et al. 1971).

During low angular accelerations, the SCCs operate in a linear push-pull fashion. For instance, during a head rotation in the horizontal plane to the right, the firing rate of afferent neurons from the right horizontal SCC increases approximately by the same magnitude as the neuronal firing rate from the left horizontal SCC decreases. This push-pull mechanism is realized in the *linear* vestibular pathways from both labyrinths to the brainstem and the cerebellum (Lasker et al. 1999; Lasker et al. 2000; Minor et al. 2009; Minor et al. 1999). Properties of the rVOR for sinusoidal head movements in the low-to-middle frequency range were successfully modelled using linear transfer functions (Paige 1983; Robinson 1981). These traditional linear models, however, could not sufficiently explain rVOR-properties observed at higher frequencies and velocities (with gain being higher than expected), as emphasized by Minor et al. (Minor et al. 1999). These authors experimentally characterized the dynamics of the rVOR at high frequencies and proposed a new model that took into account the limitations of the vestibular organs when high head accelerations are applied

(Lasker et al. 1999; Minor et al. 1999). Specifically, at high accelerations and high frequencies the change of firing rates is non-linear in that the ipsilateral increase is larger than the contralateral decrease (Lasker et al. 1999; Minor et al. 1999). The additional excitatory signal from the ipsilateral labyrinth travels within the *non-linear* pathways from both labyrinths to the brainstem and the cerebellum (Galiana et al. 2001; Halmagyi et al. 1990; Lasker et al. 1999; Lasker et al. 2000; Minor and Lasker 2009; Minor et al. 1999). This mechanism of stronger ipsilateral excitation than contralateral inhibition of ampullary hair cells during high-acceleration head rotation forms the basis of Ewald's second law (Ewald 1892). By taking both linear and non-linear pathways into account, Lasker and co-workers could experimentally reproduce Ewald's second law in canal-plugged squirrel monkeys (Lasker et al. 1999).

In humans, the existence of the non-linear vestibular pathway, and hence Ewald's second law, becomes manifest when applying the head-impulse test (Halmagyi et al. 1988) in patients with unilateral vestibular loss (Halmagyi et al. 1990). When the head is rapidly rotated to the affected side, the gain (i.e. eye-velocity divided by head-velocity) of the rVOR is lower than when the head is rotated to the unaffected side. Such deficient rVOR results in a catch-up saccade that is triggered by the visual system and that can be depicted by the clinician.

To demonstrate Ewald's second law in healthy subjects, we attempted to suppress or enhance the non-linear pathway unilaterally with caloric irrigation, and then apply rapid head rotations, since these non-linear mechanisms are elicited only for head accelerations above $\sim 100^\circ/\text{s}^2$ (Baloh et al. 1977). The non-linear pathway has been found to be especially sensitive to changes in temperature (Park et al. 2010), and in particular cold temperature both lowers the resting discharge rate and decreases the sensitivity to head velocity. The rVOR evoked by ipsilateral head impulses will thus be driven predominately by the linear pathway.

112 Conversely, warm stimulation could push the non-linear pathway into its more sensitive
113 operating range, and so should lead to an enhancement of the rVOR.

MATERIALS AND METHODS

Test subjects

Seven healthy naïve individuals (four men, three women, mean age 37 years, range 19–51 years) with no history of otologic or cochleo-vestibular disorders were recruited. The study was approved by the institutional review board of the Petz Aladár Hospital, Győr, Hungary (PAMOK Hospital Protocol number 76-1-18/2015), written informed consent was obtained from all participants. The protocol was in accordance with the ethical standards laid down in the 1964/2013 (7th revision) Declaration of Helsinki for research involving human subjects. Before the measurements the left eardrum was examined and the head-impulse test was carried out in a sitting position. All subjects had a normal eardrum and normal gain values for the horizontal rVOR.

Stimulation and measurement methods

Before warm- and cold-water caloric irrigation, baseline head-impulse testing was obtained in the plane of the lateral SCCs. Video head-impulse testing (vHIT) was carried out with ICS Impulse video goggles (GN Otometrics, Taastrup, Denmark) (MacDougall et al. 2009; Weber et al. 2009). For head-impulses, the position of the right eye was recorded with 250 samples/sec. Caloric stimulation was done with the lateral SCCs in an earth-vertical position, so the subject's head was inclined by 30° from supine, with the Reid's base line elevated 60°. Subjects were tested in a well-lit room (to ensure a small pupil) with an eye-level target at a minimum distance of 1m in front of them, which was positioned in order to allow a straight fixation-direction during the head-impulses. Caloric irrigation lasted one minute and was carried out in the left ear in all subjects and at temperatures of 24, 27, 30 and 44°C. After the one minute of irrigation, five high-acceleration head-impulses to the left and to the right (peak

head velocity $>160^{\circ}/\text{sec}$, which translates to accelerations above approximately $3000^{\circ}/\text{sec}^2$) were administered over a time interval of 30 seconds in a random order. This block of two times five impulses (both left and right) was repeated five times over 150 seconds resulting in five blocks. The gain values of impulses were averaged per block, thereby creating one data point per subject every 30 seconds. The subjects relaxed for 10 minutes between caloric irrigations. To minimize effects of possible caloric habituation due to repetitive caloric irrigation (Lidvall 1961), irrigation at different temperatures was in a different random order for each subject.

After caloric irrigation with the lateral canals in the earth-vertical position, the experiment was repeated with the lateral canals earth-horizontal (Reid's base line pitched 30 degrees down), so the head was pitched forward 30° from upright. Since in this position convectional thermal endolymphatic movements along the canal are missing, if there is any change in the rVOR, it should be caused only by direct cooling or heating of the end organ or the vestibular nerve afferents.

Analysis of caloric nystagmus

In order to assess the intensity of the caloric nystagmus during the head impulses, the nystagmus beats occurring in the time frame of individual head impulses were counted offline. The ICS software saved 750 msec of data for each head impulse, beginning 100 msec before the start of the impulse. Usually the intensity of the caloric nystagmus is characterized by the maximal slow-phase velocity of the nystagmus. With the software used in this experiment, we could not retrieve these values from the dataset. Since there is a high correlation between the slow-phase velocity and nystagmus frequency (Hinchcliffe 1968; Mulch et al. 1978), we decided to characterize the effect of irrigation using the frequency of nystagmus beats. Nonetheless, nystagmus beats could be clearly depicted and when more than

one beat was registered per trial, the nystagmus frequency (beats/second) was calculated (Figure 1). This was easily possible when the ensuing nystagmus had a direction opposite to the compensatory saccades (Figure 2 right panel). In order to differentiate caloric nystagmus beats from compensatory saccades induced by head-impulse testing when the two occurred in the same direction (Figure 2 left panel), only those nystagmus beats were counted that occurred immediately before an impulse or later than 200 msec after the impulse (i.e., at a period when vHIT-induced compensatory saccades have already stopped). In cases where caloric nystagmus quick phases and vHIT-induced compensatory saccades were expected to beat into opposite directions, this restriction was not needed. Beat frequency was only calculated, when at least two nystagmus beats could be identified unanimously. Recall that testing was done in the light; so fixation of visual targets likely suppressed the nystagmus slow phases and the beat frequency.

Head-impulse gain measurements

The head-impulse gain was calculated offline with a custom algorithm implemented in Matlab (The MathWorks, Natick, MA, USA). Due to the experimental paradigm, eye-velocity traces were prone to two distinct biases. First, caloric irrigation produces nystagmus and offsets in baseline eye velocity before the head-impulse test were recorded. Second, nystagmus beats (saccades) during the head impulses complicates gain calculation. To overcome the first bias, we removed any eye velocity offsets present in the period before the head impulse prior to gain calculation. To overcome the latter bias, velocity gain was calculated only during the ascending part of the response (i.e., before the expected onset of any compensatory saccades), over a period of 52 msec, while traces with quick phases during the critical period used for gain calculation were excluded. For our special experimental conditions, this traditional velocity gain calculation was preferred over the cumulative position algorithm (Macdougall et

al. 2013) provided by the manufacturer of the ICS Impulse Goggles. Although this method is more prone to goggle-slip artefacts around peak head acceleration, these artefacts are likely to be constant across all conditions, so that they cannot explain differences between blocks. Furthermore, it emphasizes the higher frequency response compared to the standard cumulative position algorithm provided by the manufacturer. Noteworthy, this approach is also more sensitive to lags in eye velocity relative to head velocity encountered in some traces, resulting in overall lower gains than compared to the approach provided by the manufacturer.

Statistical analysis was carried out using Graphpad® Software. Average gain values at different times after irrigation were compared using non-parametric, one-way ANOVA (Friedman test, with Dunn's multiple comparison post-test).

RESULTS

Caloric nystagmus

With the plane of the lateral SCCs positioned earth-vertical (the subject's head inclined by 30° from supine), we first confirmed the presence of caloric nystagmus evoked by irrigation of the left ear. For all subjects, Figure 1 shows the frequency and direction of nystagmus as a function of time after the end of the irrigations (average \pm 1 standard deviation, bins of 30sec duration). Water temperature above 37°C (44°C) led to caloric nystagmus in the direction of the irrigated ear (positive; slow phases: contralateral to irrigation). Thus warm-water irrigation corresponds to a long-lasting (or low-frequency) excitation of the lateral SCC on the ipsilateral side. Accordingly, water temperature below 37°C (30, 27, 24°C) led to caloric nystagmus in the contralateral direction (negative; slow phases: ipsilateral to irrigation). Over time, nystagmus frequency decreased, but was still present in most conditions 90 seconds after the end of irrigation. No caloric nystagmus was elicited when ear irrigations were performed with the plane of the lateral SCC oriented earth-horizontal (the subject's head inclined by 30° from upright; data not shown).

/* Figure 1 about here */

Figure 2 shows eye-velocity traces for a typical subject during head impulses to the left (Figure 2A) and to the right (Figure 2B) after the caloric irrigation of the left ear. Note that the head-velocity traces are mirrored for comparison with the rVOR. For cold-water (24°C) caloric irrigation of the left ear, irrigation-related nystagmus quick phases will beat to the contralateral side (i.e., rightwards). When applying head impulses to the left, the rVOR-response and the irrigation-related nystagmus will point in the same (rightward) direction (Figure 2A), whereas for head impulses to the right the leftward rVOR response and the

(rightward) irrigation-related nystagmus will point into opposite directions (Figure 2B). Note that in Figure 2 rightward eye movements are by definition always with a positive sign.

/* Figure 2 about here */

Head impulses

Gains of the rVOR elicited by head impulses were recorded before (baseline) and after ear irrigation (during caloric nystagmus). Baseline rVOR-gains for head turns to the left were somewhat smaller than those to the right (Figure 3). Most likely, this is related to the fact that the rVOR was always recorded from the right eye, and traditional velocity gains show higher values with head impulses in the direction of the eye that is recorded (Weber et al. 2008).

Figure 3 summarizes the results in all subjects. Cold-water irrigation (24°C) significantly decreased the rVOR-gain of ipsilateral head-impulse tests during the second time period (31 to 60 seconds after the end of irrigation; $p < 0.001$, one-way non-parametric ANOVA) (Figure 3A). Thus low-frequency inhibition of the lateral SCC by cold-water irrigation significantly reduced the velocity of high-frequency excitation of the same canal by ipsilateral head impulses (i.e., head impulses to the left). The rVOR-gain of contralateral (i.e., rightward) head impulses, however, was not significantly influenced by cold-water irrigation (Figure 3B). Likewise, rVOR-gains of head-impulse tests to either side were not affected by ipsilateral warm-water (44°C) caloric stimulation.

/* Figure 3 about here */

Control experiments with head-impulse tests applied to both sides after warm- and cold-water caloric stimulation with the plane of lateral SCC oriented earth-horizontal (the subject's head

250 inclined 30° from supine) did not reveal changes of rVOR-gains compared to the baseline
251 (data not shown).

DISCUSSION

The main finding of this study was that a strong cold-water caloric stimulus (24°C) had a significant inhibitory effect on the ipsilateral horizontal high-acceleration rVOR-gain, assessed with the head-impulse test. Water temperatures 7°C warmer or cooler than body temperature (44 and 30°C) were not strong enough to significantly influence the horizontal head-impulse rVOR-gain, although they were effective enough to elicit a tonic vestibular asymmetry and horizontal nystagmus.

Strong cold-water caloric irrigation offers a promising method to study the effects of at least partial acute unilateral vestibular loss in healthy human subjects. According to the model developed by Minor and Lasker, vestibular signals encoding angular head rotations are transmitted to the vestibular nuclei along two parallel afferent streams (Minor and Lasker 2009). The linear pathways from both sides contribute equally during ipsilateral and contralateral rotations, even during high accelerations. In other words, a unilateral deficit of the linear pathways would lead to the same rVOR-gain during head impulses on both sides. The lower gain on the affected side in case of unilateral peripheral vestibular loss is caused by a deficit of the non-linear pathway that normally enhances the rVOR mainly during ipsilateral horizontal head impulses. Thus, in the model of Minor and Lasker, Ewald's second law only applies for high head accelerations and is implemented by the nonlinear pathway that is driven mainly by excitatory, i.e. ipsilateral SCC input.

An alternative mechanism may be postulated based on findings that vestibular afferents change their sensitivity with temperature. Young and Anderson found that the sensitivity of regular afferents is positively correlated with temperature. Thus cold-water irrigation could decrease the sensitivity of these neurons, resulting in decreased gain for ipsilateral movements (Young et al. 1974). Park et al reported that irregular vestibular afferents show increased sensitivity with body-core temperature increases, from which we could likewise predict

decreased sensitivity with cold stimulation (Park et al. 2010). Note, however, that such a mechanism would also predict symmetrical changes in gain (increased ipsilateral gain with warm stimulation) in an earth-vertical position together with gain changes after caloric stimulation in an earth-horizontal position, which we did not observe.

A similar experiment with rotatory stimuli was recently conducted by Mantokoudis and co-workers (Mantokoudis et al. 2016). They found that moderate degrees of post-rotatory nystagmus (up to 30°/s) did not alter the rVOR-gain of horizontal head-impulses with high peak velocities. With lower peak-velocity head impulses and higher velocities of post-rotatory nystagmus (higher than 30°/s), however, the rVOR-gains increased or decreased as predicted by summation of the velocity of the post-rotatory nystagmus and the velocity of the rVOR elicited by the head impulses. These results of Mantokoudis and co-workers are, in some respect, similar to ours: With caloric stimulation of moderate temperature difference (7°C warmer or cooler than body temperature), rVOR-gains of horizontal head impulses in either direction did not change compared to baseline. With strong caloric stimulation, which was obviously only possible with below-body water temperature (24°C), we also found an effect on rVOR-gains of horizontal head impulses. Since there was no effect of caloric nystagmus when irrigating the lateral SCC in earth-horizontal position and no changes of rVOR-gains for horizontal head impulses, we conclude that the present and absent effects on the rVOR-gains in vertical position are related to the degree of ampullofugal deflection of the cupula due to thermal endolymph pressure and not by direct cooling of the ampulla and/or its afferents (Valli et al. 2002).

In the experiments of Mantokoudis and co-workers, strong post-rotatory nystagmus (nystagmus slow-phase velocity >30°/sec) increased the rVOR-gain of horizontal head impulses when both nystagmus slow-phases and head-impulse rVOR pointed in the same

direction and decreased when they were directed oppositely (Mantokoudis et al. 2016). In our study, only the strong cold-caloric stimulus (24°C) had a significant decreasing effect on the rVOR-gain of ipsilateral (i.e., leftward) horizontal head impulses, i.e., when slow-phases of caloric nystagmus and head-impulse VOR pointed in opposite directions, but there was no change of rVOR-gain during contralateral (i.e., rightward) head impulses, as illustrated in Figure 4.

/* Figure 4 about here */

The fundamental difference between the experiments of Mantokoudis and co-workers and those reported in this study is the nature of the underlying vestibular signal. While post-rotatory nystagmus is a post-deceleration velocity signal in the linear pathways coming from the SCCs of both labyrinths and is then partly integrated by the velocity-storage mechanism, caloric nystagmus is elicited by an on-going acceleration signal driven by cupular deflection in the horizontal SCC only on the side of the irrigation. The acceleration signal of a strong cold-water caloric stimulus could therefore drive the irregularly firing afferents in an area where they are relatively insensitive to high acceleration. These afferents are asymmetric in nature because they show large excitatory responses, but only small inhibitory responses (Minor and Lasker 2009). They constitute the non-linear pathway of the rVOR and implement Ewald's second law for high head accelerations. As a result of being driven into a zone less sensitive to acceleration, non-linear neurons are prevented from effectively firing during ipsilateral rapid head rotations. We hypothesize that this is the cause for the diminished rVOR-gains in the ipsilateral horizontal direction. These distinct effects of caloric irrigation on the linear and the non-linear pathway when applying head-impulses are explained in Figure 5. While, for the linear pathway, the change in the vestibular-afferent firing-rate

caused by an ipsilateral head-impulse immediately after caloric irrigation will be independent from the initial firing-rate (i.e., the “set point”), the change in firing rate in the non-linear pathway depends critically on the initial discharge rate due to the non-linear relationship between acceleration and neuronal discharge. The overall change in firing rate will therefore be lower for head impulses applied after cold-water caloric irrigation than after warm-water caloric irrigation, explaining the lower vHIT-gain (Fig 5G).

/* Figure 5 about here */

In summary, our findings emphasize the importance of the non-linear pathway when probing the vestibular organs with high-acceleration stimuli. Combining caloric irrigation and head-impulses may therefore provide helpful for further studies as a model of transient impairment of the rotational vestibulo-ocular reflex.

ACKNOWLEDGMENTS

Conflict of interest and sources of funding:

Dr. Tamás, Dr. Bockisch, Dr. Lasker, Dr. Büki, Prof. Straumann and Dr. Tarnutzer report no conflict of interest. Dr. Weber acts as an unpaid consultant and has received funding for travel from GN Otometrics.

This study was supported by Dr. Dabbous Foundation, University of Zurich, Switzerland; Swiss National Science Foundation; and Betty and David Koetser Foundation, Zurich, Switzerland.

Author contributions to the manuscript

L. T. Tamás: data collection, analysis and interpretation, critically reviewing and editing the manuscript.

K. P. Weber: data analysis and interpretation, critically reviewing and editing the manuscript.

C. J. Bockisch: interpretation of the data, assisted in writing of the manuscript.

D. Straumann: interpretation of the data, critically reviewing and editing the manuscript.

A. A. Tarnutzer: data analysis and interpretation, critically reviewing and editing the manuscript.

D. M. Lasker: interpretation of data, critically reviewing and editing the manuscript.

B. Büki: conceived of the study, data interpretation, drafted the manuscript.

All authors have read and approved the final version of the manuscript.

363 **REFERENCES**

- 364 Baloh, R. W., Honrubia, V., Konrad, H. R. (1977). Ewald's second law re-evaluated. *Acta*
365 *Otolaryngol*, 83, 475-479.
- 366 Ewald, J. R. (1892). *Physiologische Untersuchungen ueber das Endorgan des Nervus*
367 *Octavus*. Wiesbaden, Germany: J. F. Bergmann.
- 368 Fernandez, C., Goldberg, J. M. (1971). Physiology of peripheral neurons innervating
369 semicircular canals of the squirrel monkey. II. Response to sinusoidal stimulation and
370 dynamics of peripheral vestibular system. *J Neurophysiol*, 34, 661-675.
- 371 Galiana, H. L., Smith, H. L., Katsarkas, A. (2001). Modelling non-linearities in the vestibulo-
372 ocular reflex (VOR) after unilateral or bilateral loss of peripheral vestibular function.
373 *Exp Brain Res*, 137, 369-386.
- 374 Grossman, G. E., Leigh, R. J., Abel, L. A., et al. (1988). Frequency and velocity of rotational
375 head perturbations during locomotion. *Exp Brain Res*, 70, 470-476.
- 376 Halmagyi, G. M., Curthoys, I. S. (1988). A clinical sign of canal paresis. *Arch Neurol*, 45,
377 737-739.
- 378 Halmagyi, G. M., Curthoys, I. S., Cremer, P. D., et al. (1990). Head impulses after unilateral
379 vestibular deafferentation validate Ewald's second law. *J Vestib Res*, 1, 187-197.
- 380 Hinchcliffe, R. (1968). Nystagmus rate as an index of caloric test response. *Acta Otolaryngol*,
381 65, 311-315.

382 Lasker, D. M., Backous, D. D., Lysakowski, A., et al. (1999). Horizontal vestibuloocular
 383 reflex evoked by high-acceleration rotations in the squirrel monkey. II. Responses
 384 after canal plugging. *J Neurophysiol*, 82, 1271-1285.

385 Lasker, D. M., Hullar, T. E., Minor, L. B. (2000). Horizontal vestibuloocular reflex evoked by
 386 high-acceleration rotations in the squirrel monkey. III. Responses after
 387 labyrinthectomy. *J Neurophysiol*, 83, 2482-2496.

388 Lidvall, H. F. (1961). Vertigo and nystagmus responses to caloric stimuli repeated at short
 389 intervals. *Acta Otolaryngol*, 53, 33-44.

390 Macdougall, H. G., McGarvie, L. A., Halmagyi, G. M., et al. (2013). The video Head Impulse
 391 Test (vHIT) detects vertical semicircular canal dysfunction. *PLoS One*, 8, e61488.

392 MacDougall, H. G., Weber, K. P., McGarvie, L. A., et al. (2009). The video head impulse
 393 test: diagnostic accuracy in peripheral vestibulopathy. *Neurology*, 73, 1134-1141.

394 Mantokoudis, G., Tehrani, A. S., Xie, L., et al. (2016). The video head impulse test during
 395 post-rotatory nystagmus: physiology and clinical implications. *Exp Brain Res*, 234,
 396 277-286.

397 Minor, L. B., Lasker, D. M. (2009). Tonic and phasic contributions to the pathways mediating
 398 compensation and adaptation of the vestibulo-ocular reflex. *J Vestib Res*, 19, 159-170.

399 Minor, L. B., Lasker, D. M., Backous, D. D., et al. (1999). Horizontal vestibuloocular reflex
 400 evoked by high-acceleration rotations in the squirrel monkey. I. Normal responses. *J*
 401 *Neurophysiol*, 82, 1254-1270.

402 Mulch, G., Leonardy, B., Petermann, W. (1978). Which are the parameters of choice from the
 403 evaluation of caloric nystagmus? *Arch Otorhinolaryngol*, 221, 23-35.

404 Paige, G. D. (1983). Vestibuloocular reflex and its interactions with visual following
 405 mechanisms in the squirrel monkey. I. Response characteristics in normal animals. *J*
 406 *Neurophysiol*, 49, 134-151.

407 Park, H. J., Lasker, D. M., Minor, L. B. (2010). Static and dynamic discharge properties of
 408 vestibular-nerve afferents in the mouse are affected by core body temperature. *Exp*
 409 *Brain Res*, 200, 269-275.

410 Robinson, D. A. (1981). The use of control systems analysis in the neurophysiology of eye
 411 movements. *Annu Rev Neurosci*, 4, 463-503.

412 Valli, P., Buizza, A., Botta, L., et al. (2002). Convection, buoyancy or endolymph expansion:
 413 what is the actual mechanism responsible for the caloric response of semicircular
 414 canals? *J Vestib Res*, 12, 155-165.

415 Weber, K. P., Aw, S. T., Todd, M. J., et al. (2008). Inter-ocular differences of the horizontal
 416 vestibulo-ocular reflex during impulsive testing. *Prog Brain Res*, 171, 195-198.

417 Weber, K. P., MacDougall, H. G., Halmagyi, G. M., et al. (2009). Impulsive testing of
 418 semicircular-canal function using video-oculography. *Ann N Y Acad Sci*, 1164, 486-
 419 491.

420 Young, J. H., Anderson, D. J. (1974). Response patterns of primary vestibular neurons to
 421 thermal and rotational stimuli. *Brain Res*, 79, 199-212.

422

FIGURE LEGENDS

Figure 1

Nystagmus frequency after caloric irrigation (earth-vertical orientation of the lateral semicircular canal; 7 subjects; mean nystagmus frequency ± 1 standard deviation) during video-head-impulse testing. See inset for definition of symbols used.

Figure 2

Head impulses 0-30 seconds after cold (24°C) water irrigation of the left ear in a typical subject (#6), inducing a right-beating caloric nystagmus. For illustrative purposes of the relation between the beating direction of the caloric nystagmus (to the right) and the rVOR-response, we mirrored both eye and head velocity traces in panel B. Eye movements to the right were positive, and eye movements to the left were negative. Note that head velocity traces in both panels were mirrored to allow direct comparison with the eye velocity traces and that traces with quick phases during the critical period used for gain calculation were excluded. Panel A: head impulses to the left (in black) triggering an eye movement (rVOR, in green) to the right with decreased gain (compared to the head velocity trace). The star (*) indicates a quick phase to the right during the descending portion of the rVOR-response that was likely a compensatory saccade. Panel B: head impulses to the right (in black), eliciting an rVOR-response to the left (in red, i.e., with a negative sign) that did not show any decrease in gain or compensatory saccades. The right-beating fast phases depicted in panel B are consistent with caloric nystagmus beats.

Figure 3

rVOR-gains (n=7, mean ± 1 SD) for head impulses to the left (panel A) and to the right (panel

B) after cold and warm-water caloric irrigation of the left ear with the head inclined by 30° from supine (i.e., the lateral SCCs are oriented earth-vertical) over periods of 30 seconds each. Results for the different temperatures are shown separately (see inset for explanation of symbols). To improve visualization, results for the different conditions are slightly shifted along the x-axis. Significant differences are marked by brackets and stars (***: $p < 0.001$; *: $p < 0.05$), using the same color-coding as for the different measurements after caloric irrigation, i.e., green for 24° cold-water caloric irrigation and blue for 27° cold-water caloric irrigation. Non-parametric one-way ANOVA showed no significant differences between other parameters. Baseline measurements before caloric irrigation: purple diamond.

Figure 4

Hypothetical effect of cold-water caloric irrigation (24°C) of the left ear during head impulses. Red arrows represent the magnitude of changes in overall vestibular afferent activity, i.e. non-linear and linear pathway activity, during head impulses. Changes are identical in all paradigms.

Upper panel: without caloric irrigation. Columns represent the afferent discharge (spontaneous afferent activity approximately 90/sec). Head impulses to the right and left: ipsilateral afferent activity increases, the contralateral afferent discharge decreases.

Middle panel: during and after left cold caloric irrigation (blue arrows) the spontaneous afferent discharge rate decreases and the increase of the activity during ipsilateral head impulse elicits less asymmetric activity between both sides than without caloric stimulation; yellow arrow: slow phase direction of caloric nystagmus.


Lower panel: when stopped after sustained rotation to the left (counter-clockwise) (Mantokoudis et al. 2016), the endolymph movements elicit a change in the spontaneous activity as if the head would accelerate to the right (clockwise) and post-rotatory nystagmus

appears with a slow phase to the left (yellow arrow). In this case, slow phases of the post-rotatory nystagmus add or subtract to the slow-phase eye velocity during the head impulses. The difference between caloric nystagmus and post-rotatory nystagmus appears during the head impulse to the right: after caloric stimulation, there is no enhancement of the gain to the right, while there is an increase of the rVOR-gain to the right after sustained rotation.



Figure 5



Schematic relationship between changes in head acceleration (x-axis) and subsequent changes in the firing rate of the vestibular afferents (y-axis) during head impulses. The firing rate at zero head acceleration is referred to as the "set point". Regular afferents (bottom row) and irregular afferents (top row) are shown separately. Note that the relationship between firing rate and head acceleration is linear for the regular vestibular afferents (i.e., the linear pathway), while it is non-linear for the irregular vestibular afferents (i.e., the non-linear pathway). The baseline condition (no caloric irrigation before the head-impulse, panels A and D) is compared with an inhibitory condition (cold-water caloric irrigation before the head-impulse, panels B and E) and an excitatory condition (warm-water caloric irrigation before the head-impulse, panels C and F). Cold-water caloric irrigation ("inhibitory condition") shifts the set point (light grey circle) to the left (as indicated by the dashed arrow), while warm-water caloric irrigation ("excitatory condition") results in a shift to the right (as indicated by the dashed arrow). When applying a head impulse, the acceleration results in an increase in firing rate (Δ discharge rate). Due to the linear relationship, Δ discharge rate is independent from the initial firing rate. This, however, is not the case for the irregular afferents: when applying cold-water caloric irrigation, the firing rate is shifted to an area where the slope of the curve is less steep. For warm-water caloric irrigation, the firing rate is shifted to an area

496 where the slope of the curve is almost linear. Resulting overall changes in firing are shown in
497 panel G.




nystagmus
right-beating






without stimulation



left cold
stimulation



after rotatory
stimulation

